

FIRST REPORT AND PREVALENCE OF SESSILE CILIATE (*Vorticella* sp.) IN CULTURED *Clarias gariepinus* IN NIGERIA

¹OKUNADE, O. A., ²E. K. AJANI, ³J. O. ADEJINMI, AND ⁴G. A. OLADOSU

¹ Nigerian Institute for Oceanography and Marine Research, Victoria Island, Lagos State, Nigeria.

² Department of Aquaculture and Fisheries Management, University of Ibadan, Oyo State, Nigeria

³ Department of Entomology and Parasitology, University of Ibadan, Oyo State, Nigeria.

⁴ Department of Veterinary Medicine, University of Ibadan, Oyo State, Nigeria.

Corresponding author: olumideokunade@yahoo.ca 08034984980

ABSTRACT

Fish parasites commonly cause portal for secondary infections, hindering high productivity leading to economic loss in fish production. Protozoan parasites are a significant group of pathogens affecting cultured fish. The present study was carried out to investigate the prevalence of parasites infecting different stages of cultured *Clarias gariepinus* in Lagos State. A total of 487 cultured *Clarias gariepinus* were collected randomly from different culture facilities in culture systems and parasitological examinations were conducted on them. Identification keys with morphometric criteria were used to identify *Vorticella* sp from other sessile ciliates. *Vorticella* sp. was observed on the skin and intestine. It was found in all the culture systems and developmental stages. The highest prevalence for *Vorticella* sp. was found on adult skin (12.84%) and intestines (3.85 %). The mean intensity of *Vorticella* sp. was highest on fingerlings skin (23.88) and juvenile intestine (16). The mean water parameters showed a suitable range for their survival. This study is the first report of this sessile ciliate in Nigeria, indicating that more parasites are yet undiscovered in aquaculture.

Keywords: Parasitic occurrence, predilection site, *Vorticella* sp, *Clarias gariepinus*, intensity.

INTRODUCTION

The Clariid, *Clarias gariepinus* is one of the fish regularly cultured as fish food in earthen ponds or concrete tanks in Nigeria (Akinwole and Adedayo, 2014; Oluwalola, *et al.*, 2019), owing to several favorable cultural potentials (Obiekezie and Ekanem, 1995). It is a good source of protein in human diets because of its nutritive value (Alune and Andrew, 1996). Parasites and diseases have always been associated with poor production of cultured and feral fish (Subasinghe *et al.*, 2002), they often reduce the immunity of the host thereby causing ease of secondary infections resulting in nutritive devaluation and subsequent economic losses (Onyedineke *et al.*, 2010). Most often, helminths and crustacean parasites were more studied than protozoan parasites of cultured and feral fish (Lom and Dykova, 1992, Akinsanya and Otubanjo, 2006). However, more awareness has been given to parasitological studies in fish development in ponds and other fish enclosures as a result of intensification in production (Igbani, Hauwa, and Tukura, 2019). In Nigeria, several researchers have reported ecto- and endo- parasites of various fish species of catfishes (*Clarias gariepinus*, *Chrysichthys nigrodigitatus*, *Malapterurus electricus*) and Tilapias (*Oreochromis niloticus* and *Tilapia zillii*) from different aquatic ecosystems with ciliates and metazoan (Okunade *et al.*, 2018; Uhwo and Obi, 2015; Eyo, Edet, and Ekanem, 2015; Adeogun *et al.* 2014; Omeji, *et al.* 2014; Abidemi-Iromini and Eze, 2011; Omeji, Solomon and Idoga, 2011). Parasitic ciliates are pathogenic protozoans that may not be alarming in the wild but are considered to be a serious problem in aquaculture causing severe losses (Van As and

Basson, 1988; Dickerson and Clark, 1996). Many cultured fishes are vulnerable to sessile peritrichs such as *Vorticella* sp., *Apiosoma* sp., and *Ambiphrya* sp., (Abdel-Baki *et al.*, 2014; Woo and Leatherland, 2006; Viljoen and Van As 1985); *Apiosoma* sp (Sogbesan, Tukura, and Ekundayo, 2018); *Epistylis* sp (Viljoen and Van As, 1985) but the sessile ciliate; *Vorticella* sp., has not been reported in culture fish in Nigeria to the best of our knowledge. The sessile peritrich (*Vorticella* spp) found in fish are typically ectocommensals attaching to their substrates by a stalk (Viljoen and Van As, 1983), using their hosts as a living, moving substrate for convenient access to the source of food particles, organic debris and waterborne bacteria (Lom and Dykova, 1992). The species of *Vorticella* are identified using gene sequence information (Ji *et al.*, 2015; Liang *et al.*, 2018) and modern molecular methods combined with morphological characteristics (Sun *et al.*, 2013a; Wang *et al.*, 2017). *Vorticella* species of the family Vorticellidae are one of the important peritrichs commonly found in marine, freshwater, and terrestrial biotopes (Sun *et al.*, 2005; Lynn, 2008) having conical and an inverted bell-shaped body with highly contractile unbranched stalk longer than the size of the body.

They are filter feeders characterized into two forms; the trophont (possessing stalk for anchoring to the desired substratum) and a free-living telotroch (Hirst, Kita, and Dawson, 2011). *Vorticella* sp. are free-living organisms but become facultative parasites externally under adverse environmental conditions when the fishes are stressed (Basson and Van As, 2006). At high densities, sessile ciliates caused high mortality in juvenile and adult cultured fish leading to severe

economic losses globally (Van As and Basson, 1987). However, the importance of *Vorticella* sp has been reported as a target for bioengineering microfluidic devices (Ryu *et al.*, 2016), biocontrol agents for mosquitoes with a 90 % reduction in adult development (Patil, *et al.* 2016) and serves as an indicator for healthy ecosystem (Perez – Uz *et al.*, 2010).

MATERIALS AND METHODS

Sampling Techniques

Sampling was conducted from March 2017 to August 2018 and total numbers of 500 *Clarias gariepinus* were randomly collected from different culture facilities across the three Agricultural zones in Lagos State, Nigeria, and transported to the laboratory in respective jerry cans at 5 fish per container to avoid stress and mortality. A scoop net was used to harvest the fry and fingerlings while a hand net was used for juveniles and adults in concrete tanks, plastic tanks, and tarpaulin vats but dragnet for adults in earthen ponds.

Parasitological Examination

External Organs

The skin was scraped using a coverslip and the wet smear was placed on the glass slide while the operculum was opened and a small portion of the gill

$$\text{Prevalence (\%)} = \frac{\text{Number of individual host species infected with a particular parasites}}{\text{Number of hosts examined}} \times 100$$

$$\text{Intensity of infection} = \frac{\text{Total number of particular parasites species in a sample of a host species}}{\text{Number of infected individuals of the host species in the sample}}$$

Water Quality Measurement

The pH, water temperature, and dissolved oxygen were recorded using Hanna digital meters at the sampling site. Transparency was measured with Secchi disc and titrimetric method using respective Hanna reagents for ammonia, nitrite, iron, alkalinity, and hardness according to (APHA, 2005).

RESULTS

Occurrence and prevalence of *Vorticella* sp

The sessile ciliate species, *Vorticella* sp are characterized into two forms; the trophont (possessing stalk for anchoring to the desired substratum) and a free-living telotroch (Plate 1a, 1b, and 1c) were identified on the skin and intestine of developmental stages of cultured *Clarias gariepinus* (Table 1). The parasitological examination revealed the occurrence of *Vorticella* sp., for the first time in culture systems in Nigeria. *Vorticella* sp. were found in all the culture facilities used for rearing and

was cut and placed on the glass slide with a drop of distilled water and viewed under Olympus microscope at high power magnification (×40) for protozoan parasites (Hanish, 2010; Ekanem *et al.* 2011) and identified using appropriate keys (Viljoen and Van As, 1983; Lom and Dykova, 1992; Jayasree *et al.*, 2001).

Internal Organs

The internal organs observed were the intestine, liver, and trunk kidney. The abdomens of the fish (excluding the fry) were incised with scissors and a scalpel to open the visceral to locate the desired organs. The intestine was longitudinally incised and the waste was removed to expose the intestinal wall which was scraped and the content was smeared on the glass slide for parasitic examination. The quantum size of the liver and trunk kidney was harvested each using scissors and placed on the glass slide, slightly masticated and compressed with another glass slide, and viewed under the microscope.

Prevalence of Parasites and Mean Intensity

The prevalence (%) of the parasite and mean intensity levels were determined using the method of Bush *et al.* (1997).

developmental stages. They were found on the skin of 75 – 100 % of the fish stages harvested in all culture facilities except plastic tanks with 25% occurrence (Table 1).

Vorticella sp. occurred in intestines of fingerlings and juveniles reared in plastic and tarpaulin vat respectively. The prevalence of *Vorticella* spp on fish skin was highest in adults (12.84%) followed by fry (9.41%) and least in juveniles (3.54%). On the intestine, the highest prevalence of *Vorticella* sp was found in adults (3.85) (Table 2).

The intensity of *Vorticella* sp on the developmental stage of *Clarias gariepinus*

The intensity of *Vorticella* sp on the skin of *Clarias gariepinus* was highest on fingerlings (23.88) followed by juveniles (14) whereas juveniles had the highest intensity (16) in the intestine examined (Table 2).




Plate 1a: *Vorticella* sp x 400




Plate 1b: *Vorticella* sp x 400



Plate 1c. Telotroch of *Vorticella* sp x 100

Plate 1a and 1b : C – Cilia, F – Food vacuole, SC – Scopula, ST – stalk, PL – Peristomial lip, PD – Peristomial disc, Z - Zooid

Table 1: Occurrence of *Vorticella* sp on the skin and in the intestine of developmental stages of *Clarias gariepinus* reared in different culture facilities

| Culture Facilities | Developmental Stages | Site of location <i>Vorticella</i> sp. | | | |
|--------------------|----------------------|--|------|-----------|--|
| | | | Skin | Intestine | |
| Earthen ponds | Fry | - | - | - | |
| | Fingerlings | | + | - | |
| | Juveniles | | + | - | |
| Concrete tanks | Fry | + | - | - | |
| | Fingerlings | | + | - | |
| | Juveniles | | + | - | |
| Plastic tanks | Fry | + | - | - | |
| | Fingerlings | | - | + | |
| | Juveniles | | - | - | |
| Tarpaulin vats | Fry | + | - | - | |
| | Fingerlings | | + | - | |
| | Juveniles | | + | + | |
| | Adults | | - | - | |

Table 2: Prevalence and intensity of *Vorticella* sp. on the skin and in the intestine of developmental stages of cultured *Clarias gariepinus* in Lagos State.

| Developmental Stage | Numbers of fish examined | NFI | Skin | | NFI | Intestine | |
|---------------------|--------------------------|-----|----------------|-----------|-----|----------------|-----------|
| | | | Prevalence (%) | Intensity | | Prevalence (%) | Intensity |
| Fry | 85 | 8 | 9.41 | 9.25 | 0 | 0 | 0 |
| Fingerlings | 95 | 8 | 8.42 | 23.88 | 1 | 1.05 | 7 |
| Juveniles | 198 | 7 | 3.54 | 14 | 2 | 1.01 | 16 |
| Adults | 109 | 14 | 12.84 | 10 | 4 | 3.85 | 8.5 |

NFI – Number of infected fish

Water Quality Parameters

The water parameters recorded in culture systems for *Vorticella* sp. in this study were shown in table 3. The range of water parameters under which *Vorticella* sp thrived were pH (6 – 8), water

temperature (29 – 33 °C), dissolved oxygen (1 - 8mg^l⁻¹), ammonia (1 – 3.3mg^l⁻¹), nitrite (0.06 – 1.30 mg^l⁻¹), iron (0 – 1.40 mg^l⁻¹), alkalinity (28 – 128 mg^l⁻¹), hardness (90 – 760 mg^l⁻¹) and transparency (22 – 50 cm).

Table 3: Water quality parameters in culture tanks of *Clarias gariepinus* with *Vorticella* sp.

| Water Parameters | <i>Vorticella</i> sp |
|--|----------------------|
| Water temperature (°C) | 30.77 ± 1.58 |
| pH | 6.99 ± 0.81 |
| Dissolve Oxygen (mg ^l ⁻¹) | 4.30 ± 3.01 |
| Ammonia (mg ^l ⁻¹) | 2.58 ± 0.64 |
| Nitrite (mg ^l ⁻¹) | 0.64 ± 0.58 |
| Iron (mg ^l ⁻¹) | 0.54 ± 0.85 |
| Alkalinity (mg ^l ⁻¹) | 78.18 ± 49.69 |
| Hardness (mg ^l ⁻¹) | 433.64 ± 341.78 |
| Transparency (cm) | 35.76 ± 13.40 |

DISCUSSIONS

Vorticella sp was found on the skin of cultured *Clarias gariepinus* in this study conforming to the findings of several researchers on different fish species such as cultured *Oreochromis niloticus* (Abdel – Baki *et al.*, 2014), wild *Sarotherodon galilaeus* (Reda, 2011), *Cyprinus carpio* and *Ctenopharyngodon idella* (Guguloth *et al.*, 2013) contrary to the reports on *Vorticella* sp., found in the gills of *Clarias gariepinus* (El-Tantawy and El-Sherbiny 2010); *C. carpio* and *C. idella* (Dash *et al.*, 2015) and skin and gills of *Planiliza abu* (Al-Musawi and Al-Rubaie, 2018). The *Vorticella* sp., found on all the stages implies that they were found in a hatchery and grow-out. The presence of parasites on the skin may be due to their first contact in the culture environment as well as large surface to search for food similar to the report on *Vorticella* sp., attaching to the fin of flounder, *Platichthys flesus* L (Öztürk and Özer, 2010). The presence of *Vorticella* sp in the intestine may be accidental through ingestion since the intestine recorded minimal values in the study which may agree with the free-living nature of the parasite by preferring the external organs for attachment for ease of accessing the source of food particles such as waterborne bacteria and organic debris (Lom and Dyková, 1992). The prevalence of *Vorticella* sp., in all the developmental stages, ranged from 3 – 13 % which has similar trends of 8 – 11 % and 9.25 – 15 % in *Cyprino carpio* and *Ctenopharyngodon idella*

respectively (Gadadhar, Debolina, and Raghu, 2015). The sessile ciliate was able to withstand stress conditions range of water parameters especially dissolved oxygen (1 – 8 mg^l⁻¹), ammonia (1 – 3.3 mg^l⁻¹), nitrite (0.06 – 1.30 mg^l⁻¹) hardness (90 – 780 mg^l⁻¹). The implication is that *Vorticella* sp will comfortably adapt and survive in a suitable and probably stressed culture environment as long as the hosts are alive.

CONCLUSION

The report of *Vorticella* sp represents the first record of the parasites in *Clarias gariepinus* and culture fish in Nigeria. There are many protozoans and metazoans yet to be discovered and reported for aquaculture interest to improve the management protocol in fish health for better yield in fish produce.

REFERENCE

- Abdel-Baki, A. S., Gewik, M. M. and Al-Quraishy, S. (2014). First records of *Ambiphrya* and *Vorticella* spp. (Protozoa, Ciliophora) in cultured Nile tilapia (*Oreochromis niloticus*) in the central region of Saudi Arabia. *Saudi Journal of Biological Sciences*; 21: 520 – 523.
- Abdemi-Iromini, A.O. and Eze, R.N. (2011). Comparative assessment of parasite infestation of Tilapia in natural and cultured environments. *Proceedings of the Ninth International Symposium on Tilapia in aquaculture*. (Eds.)

- Liping, L. and Fitzsimmons, K. Shanghai Ocean University, Shanghai, China, 22–24.
- Adeogun, O.A., Oladosu, G.A., Akinwale, M.M.A., Okunade, O.A., Akintayo, I.A., Idika, N., Adeiga, A.A., Ezeugwu, S.M.C., Afocha, E.E, Peters, O.S. and Odusanya, F.A. (2014). Identification, Distribution, and Prevalence of Ecto-parasites Associated with cultured fish in Ogun State, Nigeria. *Journal of Fisheries and Aquatic Science*; 9(5):413 - 418
- Akinsanya, B. and Otubanjo, O. A. (2006). Helminth parasites of *Clarias gariepinus* (clariidae) in Lekki lagoon, Lagos, Nigeria. *Revista de Biologia Tropical*, 54(1):93-99.
- Akinwale, A. O. and Adedayo, P. A. 2014. Influence of tanks liner material on water quality and growth of *Clarias gariepinus*. *Journal of Agriculture and Social Research*, 14(1): 85 – 90.
- Al-Musawi, A. K and Al – Rubaie, A. L. (2018). The First Record of Two Species of Protozoa: *Vorticella globularia*
- Alune E and Andrew G. (1996). *Fishes* (8th ed.). Cambridge, University Press, London, UK. Pp 225.
- APHA. (2005). American Public Health Association, *Standard Methods for the Examination of Water and Wastewater*, American Public Health Association, the American Water Works Association, and the Water Environment Federation, Washington, DC, USA, 20th edition, 2005.
- Basson. L. and Van As, J. G. (2006). Trichodinidae and other ciliophorans (Phylum Ciliophora). In: Woo PTK. *Fish diseases and disorders*. 2nd ed. vol. 1. Protozoan and Metazoan infections p. 154-182. Cab International. London.
- Bush, A.O., Lafferty, K.D., Lotz, J.M. and Shostak, A.W. (1997). Parasitology meets ecology on its own terms: Margolis, L., Esch, G.W., Holmes, J.C., Kuris, A.M and Schad, G.A. (1982) revisited. *Journal of Parasitology*; 83: 575-583.
- Dash, G., Majumder, D. and Raghu, R. K. (2015). Seasonal distribution of parasites in freshwater exotic carps of West Bengal, India. *Indian Journal of Animal Research*; 49 (1): 95-102.
- Dickerson, H.W. and Clark, T.G., (1996). Immune response of fishes to ciliates. *Annual Review of Fish Disease*; 6, 107-120.
- Ekanem, A.P., Eyo, V.O. and Sampson, A.F. (2011). Parasites of landed fish from Great Kwa River, Calabar, Cross River State, Nigeria. *International Journal of Fisheries and Aquaculture*; 3(12):225-230.
- El-Tantawy, S.A.M. and El-Sherbiny, H.A.E. (2010). Some Protozoan Parasites Infecting Catfish *Clarias gariepinus* Inhabiting Nile Delta Water of the River Nile, Dakahlia Province, Egypt. *Journal of American Science*; 6(9), 676 – 696
- Eyo, V. O., Edet, T. A. and Ekanem, A. P. (2015). Monogean parasites of the African catfish *Clarias gariepinus* from two fish farms in Calabar, Cross River State, Nigeria. *Journal of Coastal Life Medicine*, 3(6):433-437. DOI: 10.12980/JCLM.3.2015J5-9. www.jclmm.com.
- Fernandez-Leborans, G. (2004). Protozoan epibionts on *Mysis relicta* Loven, 1862 (Crustacea, Mysidacea) from Lake Lušiai (Lithuania). *Acta Zoologica* (Stockholm) 85: 101-112.
- Gadadhar, D., Debolina, M, and Raghu, R. K. (2015) Seasonal distribution of parasites in freshwater exotic carps of West Bengal, India. *Indian Journal of Animal Research*; 49 (1): 95-102
- Guguloth, B., Ramudu., K. R., Subbaiah., K. and Rajesh, S. C. (2013) Prevalence of parasite diseases in carp in Bheries of West Bengal, India. *International Journal of Bio-Resource and Stress Management*; 4 (3): 468-474.
- Hamish, D. R. (2010) *Fish Disease Manual*. Vet-Aqua International, Oranmore, Co. Galway, Ireland.
- Hirst, M. B., Kita, K. N. and Dawson, S. C. 2011. Uncultivated Microbial Eukaryotic Diversity: A Method to Link ssu rRNA Gene Sequences with Morphology. *PLoS One* 6 (12), e28158 <https://doi.org/10.1371/journal.pone.0028158>
- Igbani, F., Hauwa, O. S. and Tukura, E. E. (2019) A review of the parasites of catfishes and tilapias in the wild and homestead ponds in Nigeria. *International Journal of Fisheries and Aquatic Studies*; 7(5): 307-310
- Jayasree, L., Janakiram, P. and Madhavi, R. (2001) Epibionts and parasites of *Macrobrachium rosenbergii* and *Metapenaeus dobsoni* from Gosthani estuary. *Journal of Natural History*; 35: 157-167.
- Ji, D.D., Kim, J. H., Shazib, S.U.A., Sun, P., Li, L.Q. and Shin, M. K., (2015). Two new species of Zoothamnium (Ciliophora, peritrichia) from Korea, with new observations of *Z. parahentscheli*. *Journal of Eukaryotic Microbiology*; 62:505–518.
- Liang, Z., Shen, Z., Zhang, Y., Ji, D., Li, J., Warren, A., and Lin, X. (2018). Morphology and phylogeny of four new *Vorticella* species (Ciliophora: peritrichia) from coastal waters of southern China. *Journal of Eukaryotic Microbiology* 0, 1–14. DOI: 10.1111/jeu.12668
- Lom, J. and Dykova L. (1992) *Protozoan Parasites of Fish* (Developments in Aquaculture and Fisheries Science, 26). Elsevier, Science Publishers, Amsterdam. 315 Pp
- Lynn, D. H. (2008). *The Ciliated Protozoa. Characterization, Classification, and Guide to the Literature*. 3rd ed. Springer, Dordrecht.
- Obiekezie, A. & Ekanem, D. (1995) Experimental infection of *Heterobranchus longifilis* (Teleostei, Clariidae) with *Trichodina maritinkae* (Ciliophora, Peritrichida). *Aquatic Living Resources*, 8 (4):439-443.
- Okunade, O.A., Oladosu, G. A., Adeogun, O. A., Akinwale, M.M.A., Akintayo, I. A., Peters, O. S.,

- and Odusanya, A. F. (2018). Prevalence of monogenea trematodes in different developmental stages of *Clarias gariepinus* in Delta State, Nigeria. *Nigerian Journal of Fisheries* 15 (2);1449 – 1452
- Oluwalola, O.I., Fagbenro, O.A. and Adebayo, O.T. 2019. Growth and Economic Performances of African Catfish, *Clarias gariepinus* Reared under Different Culture Facilities. *African Journal of Resources Management Fisheries and Aquatic*; 4: 75 – 84. ISSN: 2672-4197 (Prints) ISSN: 2672-4200 (Online).
- Omeji, S., Solomon, S.G. and Idoga, E.S. (2011) A Comparative Study of the Common Protozoan Parasites of *Clarias gariepinus* from the Wild and Cultured Environments in Benue State, Nigeria. Hindawi Publishing Corporation. *Journal of Parasitology Research*, Article ID 916489, 8p DOI: 10.11555/2011/916489.
- Omeji, S., Tihamiyu, L.O., Annune, P.A. and Solomon, S.G. (2014) Ecto and Intestinal Parasites of *Malapterus electricus* from Upper River Benue, *Journal of Global Biosciences* 3(6); 895-903
- Onyedineke NE, Obi U, Ofoegbu PU, Ukogo I. (2010) Helminth parasite of some Freshwater fish from River Niger at Illushi, Edo State, *Nigerian Journal of Animal Science*; 6(3):16-21
- Öztürk, T. and Özer, A. (2010). The occurrence of epizotic ciliates (Protozoa: Ciliophora) of the juvenile flounder, *Platichthys flesus* L., 1758, from Sarikum Lagoon Lake (Sinop, Turkey). *Turkish Journal of Zoology*; 34: 321-332
- Patil, C. D., Narkhede, C. P., Suryawanshi, R. K. and Patil, S. V. (2016). *Vorticella* sp: Prospective Mosquito Biocontrol Agent. *Journal of Arthropod-Borne Diseases*; 10:602–607.
- Pérez-Uz, B., Arregui, L., Calvo, P., Salvadó, H., Fernández, N., Rodríguez, E., Zornoza, A. and Serrano, S. (2010). Assessment of plausible bio-indicators for plant performance in advanced wastewater treatment systems. *Water Resources* 44:5059–5069.
- Reda, E. S. A. (2011) A Review of some ecto- and endo- protozoan parasites infecting *Sarotherodon Galilaeus* and *Tilapia zillii* from Damietta branch of River Nile, Egypt. *Journal of American Science*, 7:362-373.
- Ryu, S., Pepper, R. E., Nagai, M. and France, D. C. (2017). *Vorticella*: A protozoan for bio-inspired engineering. *Micromachines* 8:1–25.
- Sogbesan O. A., Tukura E. E. and Ekundayo, T. M. (2018). Parasitic infection and prevalence in *Clarias gariepinus* in Lake Gerio, Yola, Adamawa state. *MOJ Anatomy and Physiology*; 5(6):376–381
- Subasinghe, R.P. and Philips, M. J. (2002). Aquatic Animal Health Management; Opportunities and Challenges for Rural, Small-Scale Aquaculture and Enhanced Fisheries Development; Workshop Introductory Remarks. In Primary Aquatic Animal Health Care in Rural, Small-Scale Aquaculture Development, ed. J.R. Arthur, M.J. Philips, Subasinghe, M.B. Reantaso, and I. H. MacRae, 1-5. FAO Fisheries Technical Paper 406, FAO, Rome.
- Sun, P., Song, W., Clamp, J. and Al-Rasheid, K. A. S. (2005). Taxonomic characterization of *Vorticella fusca* Precht, 1935 and *Vorticella parapulchella* n.sp., two marine peritrichs (Ciliophora, Oligohymenophorea) from China. *Journal of Eukaryotic Microbiology*, 53: 348–357
- Sun, P., Clamp, J.C., Xu, D., Huang, B., Shin, M.K., and Turner, F., (2013a). An ITS-based phylogenetic framework for the genus *Vorticella*: finding the molecular and morphological gaps in a taxonomically difficult group. Proceedings of the Royal Society – Biological Sciences 280, 20131177.
- Tassaduqe, K., Ali, M., Salam, A., Latif, M. and Zahra, T. (2003). Study of the seasonal variations in the physicochemical and biological aspects of Indus River Pakistan. *Pakistan Journal of Biological Sciences*. 6: 1795-1801.
- Uneke, B. I., Uhuo, C. and Obi, C. (2015). Protozoan Parasites of *Chrysichthys nigrodigitatus* (Lacepede: 1803) in the Mid-Cross River Flood System, South Eastern Nigeria. *American Journal of Microbiology and Biotechnology*, 2(4):51-56.
- Van As, J. G. and Basson, L. (1987). Host specificity of trichodinid ectoparasites of freshwater fish. *Parasitology Today* 3: 88-90.
- Van As, J.G. and Basson, L., (1988). The incidence and control of fish ectoparasitic protozoa in South Africa. Technical Communication, No. 211 Republic of South Africa.
- Viljoen, S. and Van As, J.G. (1983) A taxonomic study of sessile peritrichians of a small impoundment with notes on their substrate preferences. *Journal of the Limnological Society of Southern Africa*; 9: 33 – 42.
- Viljoen, S. and J.G. van As., J. G. (1985) Sessile peritrichs (Ciliophora: Peritricha) from freshwater fish in the Transvaal, South Africa. *South African Journal of Zoology*, 20:3, 79-96, DOI: 10.1080/02541858.1985.11447920
- Wang, Z., Zhou, T., Guo, Q., Gu, Z. (2017). Description of a new freshwater ciliate *Epistylis wuhanensis* n. sp. (Ciliophora, peritrichia) from China, with a focus on phylogenetic relationships within family epistylididae. *Journal of Eukaryotic Microbiology*; 64, 394–406.
- Woo, P.T.K. and Leatherland, J. F. (2006). Fish Diseases and Disorders, second ed. CABI Pub., Wallingford, UK; Cambridge, MA.